



PRAIRIE VISTA VETERINARY HOSPITAL

Sheep and Goat Vaccines

Kids:

Covexin 8 (Clostridial with tetanus) or other Clostridium C,D and Tetanus

- Give at 6-8 weeks of age
- Booster in 2-4 weeks

Pneumonia Vaccines =

Mannheimia Haemolytica, Pasteurella Multocida

- 8-12 weeks of age
- Booster in 2-4 weeks

Does:

Campylobacter fetus (Vibriotic abortion, vibriosis)

- 1st year 1 shot just prior to or following breeding
 - o booster in 3rd Trimester or 4 months later
- In subsequent years give one shot just prior to or following breeding

Chlamydia psittici (enzootic abortion)

- Two doses are recommended. Give the first dose at least 60 days before ewes are exposed to rams, followed by a second dose 30 days later. Revaccinate annually.

Covexin 8 (Clostridial with tetanus) or other Clostridium C,D and Tetanus

- booster yearly 1 month prior to giving birth

Pneumonia Vaccines =Mannheimia Haemolytica, Pasteurella Multocida

- booster yearly 1 month prior to giving birth

Bucks:

Covexin 8 (Clostridial with tetanus) or other Clostridium C,D and Tetanus

- booster yearly with ewes

Pneumonia Vaccines =Mannheimia Haemolytica, Pasteurella Multocida

- booster yearly with ewes